arXiv:physics/0111156v1 [physics.plasm-ph] 21 Nov 2001

Kinetic temperature gradient driven modes in inhomogeneous plasmas

A. I. Smolyakov

Department of Fusion Research, Japan Atomic Energy Research Institute,
Naka, Japan
and Department of Physics and Engineering Physics, University of
Saskatchewan, Saskatoon, S7N5E2 Canada

M. Yagi

Research Institute for Applied Mechanics, Kyushu University, Japan

Y. Kishimoto

Department of Fusion Research, Japan Atomic Energy Research Institute, Naka,
Japan

(February 2, 2008)

Abstract

New unstable temperature gradient driven modes in an inhomogeneous plasma are identified. These modes represent transient $\omega \simeq k_{\|} v_{th}^{(e,i)}$ sound oscillations in magnetized plasma that are kinetically destabilized via Landau interactions. Electron and ion sound branches are unstable for large values of the Larmor radius parameter $k_{\perp}\rho_{e,i} \gg 1$, respectively. The instability occurs due to a specific plasma response that significantly deviates from Boltzmann distribution in the region $k_{\perp}\rho_{i,e} \gg 1$. Pacs: 52.35 Kt, 52.35 Qz

Small scale instabilities driven by temperature gradients are believed to be responsible for particle and energy transport in a tokamak. The ion temperature gradient driven turbulence has been identified [1] as a primary cause of ion thermal transport. The electron temperature gradient driven modes are believed to be largely responsible for the electron transport [2]. Both types of modes, the ion temperature gradient (ITG) and electron temperature gradient (ETG), have been extensively studied over last decades [3–9] In basic slab geometry these modes are essentially sound waves destabilized by coupling to pressure fluctuations [3–7]. In toroidal geometry the modes are driven by the unfavourable magnetic curvature rather than by acoustic oscillations, though the mode may remain essentially slab-like in the region of the weak/negative shear [19,20]. In this work we report on a new mechanism of the destabilization of the temperature gradient driven modes. The mechanism is essentially related to the wave-particle Landau interaction. However, the most important element is a newly found regime of plasma response for large values of the Larmor radius parameter, $k_\perp^2\rho_\alpha^2 \gg 1$, $\alpha = (e, i)$. In this regime plasma response in essential way deviates from usually assumed Boltzmann distribution for $k_\perp^2\rho_\alpha^2 \gg 1$. As is shown below modified plasma response occurs for transient (acoustic type) modes whose eigen-frequency $\omega \simeq k_\| v_{th\alpha}$ does not grow with increase of $k_\perp^2\rho_\alpha^2$, but rather remains constant. As a result two new branches of the temperature gradient driven modes occur for $k_\perp^2\rho_i^2 \gg 1$ and $k_\perp^2\rho_e^2 \gg 1$. We call them ion and electron kinetic temperature gradient (KTG) modes, respectively.

We consider shearless dispersion equation in a local approximation. Within the local theory the parallel velocity and density perturbations for each species are given by standard expressions (see e.g. [12,17])

$$\frac{\widetilde{v}_{\alpha\|}}{v_{th\alpha}} = -\frac{e_\alpha\widehat{\phi}}{T_\alpha} s_\alpha D_\alpha, \tag{1}$$

$$\frac{n_\alpha}{n_0} = -\frac{e_\alpha\phi}{T_\alpha} l_\alpha - \frac{e_\alpha\widehat{\phi}}{T_\alpha} D_\alpha. \tag{2}$$

Here

$$\begin{aligned} D_\alpha = & \left(1 - \frac{\omega_{n\alpha}}{\omega}\right)(1 + s_\alpha Z(s_\alpha))\,\Gamma_0(b_\alpha) + \frac{\omega_{T\alpha}}{\omega} s_\alpha \left(\frac{1}{2}Z(s_\alpha) - s_\alpha - s_\alpha^2 Z(s_\alpha)\right)\Gamma_0(b_\alpha) \\ & + \frac{\omega_{T\alpha}}{\omega}(1 + s_\alpha Z(s_\alpha))\,(\Gamma_0(b_\alpha) - \Gamma_1(b_\alpha))\, b_\alpha, \end{aligned} \tag{3}$$

and

$$l_\alpha = 1 - \left(1 - \frac{\omega_{n\alpha}}{\omega}\right)\Gamma_0(b_\alpha) - \frac{\omega_{T\alpha}}{\omega}\left(\Gamma_0(b_\alpha) - \Gamma_1(b_\alpha)\right) b_\alpha. \tag{4}$$

Various plasma parameters are defined as follows: $\omega_{n\alpha} = -k_y cT_\alpha/e_\alpha B_0 L_n$, $\omega_{T\alpha} = -k_y cT_\alpha/e_\alpha B_0 L_{T\alpha}$, $L_n^{-1} = -n_0^{-1}\partial n_0/\partial x$, $L_{T\alpha} = -T_\alpha^{-1}\partial T_\alpha/\partial x$, $s_\alpha = \omega/k_\parallel v_{th\alpha}$, $b_\alpha = k_\perp^2\rho_\alpha^2/2$, $v_{tha}^2 = 2T_\alpha/m_\alpha$, $\rho_\alpha = v_{tha}m_\alpha c/(e_\alpha B_0)$; $\Gamma_{0,1}(b) = I_{0,1}\exp(-b)$, and $Z(s)$ is the standard plasma dispersion function. We have introduced auxiliary potential $\widehat{\phi} = \phi - \omega/(k_\parallel c)A$, where ϕ is the electrostatic potential and A is the magnetic vector potential.

The vector potential A can be found from the Ampere's law

$$J_\parallel = en(v_{i\parallel} - v_{e\parallel}) = -\frac{c}{4\pi}\nabla_\perp^2 A. \tag{5}$$

By using (1) and (5) one obtains

$$A = -\frac{2s_e^2\left(D_e\tau + D_i\right)}{k_\perp^2\delta^2 - 2s_e^2\left(D_e\tau + D_i\right)}\frac{k_\parallel c}{\omega}\phi, \tag{6}$$

and

$$\widehat{\phi} = \phi\frac{k_\perp^2 c^2/\omega_{pe}^2}{k_\perp^2\delta^2 - 2s_e^2\left(D_e\tau + D_i\right)}, \tag{7}$$

for the parallel potential $\widehat{\phi}$. Here $\tau = T_e/T_i$, and $\delta^2 = c^2/\omega_{pe}^2$.

By using the Poisson equation,

$$-\nabla_\perp^2\phi = 4\pi e(n_i - n_e), \tag{8}$$

and expressions (2) and (7) we obtain general local dispersion relation

$$\begin{aligned} &k_\perp^2\delta^2\left(l_i\tau + l_e + D_i\tau + D_e\right) - 2s_e^2\left(D_i\tau + D_e\right)\left(l_i\tau + l_e\right) \\ &= -k_\perp^2\lambda_D^2\left(k_\perp^2\delta^2 - 2s_e^2\left(D_i\tau + D_e\right)\right), \end{aligned} \tag{9}$$

where λ_D^2 is the Debye lenghth, $\lambda_D^2 = T_e/(4\pi n_0 e^2)$. This general dispersion equation describes both ITG and ETG modes as well as new kinetic temperature gradient driven modes.

To illustrate existence of a new unstable mode we plot a solution of the dispersion equation (9) as a function of the normalized ion Larmor radius parameter ρ_i/L_n. The following parameters are fixed $k_y\rho_i = \sqrt{2}\times 0.3$, $k_x\rho_i = \sqrt{2}\times 0.1$, $\beta = 2\times 10^{-4}$, $\rho_i/L_{Ti} = \sqrt{2}\times 0.1$, $\rho_i/L_{Te} = \sqrt{2}\times 0.01$, $k_\parallel\rho_i = \sqrt{2}\times 0.002$, $\tau = 1$, We have chosen the above plasma parameters to be exactly the same as in Ref. 17. The plasma pressure parameter β is defined as $\beta = 8\pi n_0 T_i/B_0^2$. [Note that our definition of $v_{th\alpha}^2$ and β differ differ from those in Ref. 17 by a factor of 2.] Figures 1a and 1b show the real ω_r and imaginary ω_i parts of the eigen frequency, respectively. It is already evident from Figs 1a and 1b that there are exist two distinct eigenmodes: in Fig. 1b the left curve is the unstable ITG branch and the right curve is the unstable drift mode branch [17] (in terminology of Ref. 17). Our results in Figs 1a and 1b are in complete agreement with Ref. 17.

To clarify the nature of the unstable branch we solve the dispersion equation as a function of the $k_y\rho_i$ parameter for fixed $\beta = 2\times 10^{-4}$, $\rho_i/L_n = \sqrt{2}\times 10^{-2}$, $\rho_i/L_{Ti} = \sqrt{2}\times 10^{-1}$, $\rho_i/L_{Te} = \sqrt{2}\times 10^{-1}$, $k_x\rho_i = \sqrt{2}\times 10^{-1}$, $\tau = 1$. We also take $\lambda_D = 0$ for this case. The mode growth rate and mode frequency normalized to $k_\parallel v_{the}$ are shown in Figs. 2a and 2b, respectively. Two new unstable branches exist in the regions $k_y\rho_i \geq 1$ and $k_y\rho_e \geq 1$. The standard ITG mode has a peak growth rate around $k_y\rho_i \simeq 1$, and the standard ETG mode has a growth rate peaked at $k_y\rho_i \simeq 40$ (corresponding to $k_y\rho_e \simeq 1$). A new ion mode (ion KTG) has a maximum growth rate around $k_y\rho_i \simeq 5$, and a new electron mode (electron KTG) saturates to a constant growth rate quickly for large $k_y\rho_i > 200$ ($k_y\rho_e > 5$). It is essential to note that new modes exist in the regions of large Larmor radius parameters (respectively, $k_y\rho_i$ for the ion KTG, and $k_y\rho_e$ for the electron ETG). As we show below, kinetic temperature gradient modes occur due to a peculiar behaviour of the plasma response in the region of large $k_y\rho_\alpha$. Let us consider this response in more details.

A standard notion is that for large values of the Larmor radius parameter, $k_y\rho_\alpha > 1$, the density response of the respective plasma component is Boltzmann due to decaying asymptotics of $\Gamma_{0,1}(b_\alpha) \sim 1/\sqrt{b_a}$ for large b_α, so that we have from (3-4) $D_\alpha \ll 1, l_\alpha \simeq 1$. This, in fact, implicitly assumes that the ratio of $\omega_{n\alpha}/\omega$ is finite for large $k_y\rho_\alpha$. In turn, this requires that the mode eigen-frequency increases with $k_y\rho_\alpha$ (linearly or faster). The latter is definitely true for drift wave type modes, where $\omega \sim \omega_{*\alpha}$. However, temperature gradient driven modes, are basically sound waves whose frequency is of the order of the transient (or sound) frequency, $k_\parallel v_{th\alpha}$. If the parameter $k_\parallel v_{th\alpha}$ remains constant (so ω is approximately constant), the ratio $\omega_{n\alpha}/\omega$ will be increasing with $k_y\rho_\alpha$ that may compensate for the decaying $1/\sqrt{b_a}$ factor from $\Gamma_{0,1}(b_\alpha)$. Then, for large $k_y\rho_\alpha \gg 1$ we may write in the leading order

$$\frac{\omega_{n\alpha}}{\omega}\Gamma_0(b_\alpha) = \frac{1}{2\sqrt{\pi}}\frac{e_\alpha}{e}\frac{v_{th\alpha}}{\omega L_n}, \tag{10}$$

and

$$\frac{\omega_{T\alpha}}{\omega}\left(\Gamma_0(b_\alpha) - \Gamma_1(b_\alpha)\right) b_\alpha = \frac{1}{4\sqrt{\pi}}\frac{e_\alpha}{e}\frac{v_{th\alpha}}{\omega L_{T\alpha}}. \tag{11}$$

By using these asymptotics for the electron KTG mode in the region $k_y\rho_e \gg 1$ and $k_y\rho_i \gg 1$ we obtain the following response functions

$$l_i = 1 - \frac{1}{2\sqrt{\pi}}\frac{v_{ti}}{\omega L_n} + \frac{1}{4\sqrt{\pi}}\frac{v_{ti}}{\omega L_{Ti}}, \tag{12}$$

$$l_e = 1 + \frac{1}{2\sqrt{\pi}}\frac{v_{te}}{\omega L_n} - \frac{1}{4\sqrt{\pi}}\frac{v_{te}}{\omega L_{Te}}, \tag{13}$$

$$D_i = -\frac{1}{4\sqrt{\pi}}\frac{v_{ti}}{\omega L_n}\frac{1}{s_i^2} + \frac{1}{4\sqrt{\pi}}\frac{v_{ti}}{\omega L_{Ti}}\frac{1}{s_i^2}, \tag{14}$$

$$\begin{aligned} D_e = &-\frac{1}{2\sqrt{\pi}}\frac{v_{te}}{\omega L_n}\left(1 + s_e Z(s_e)\right) + \frac{1}{2\sqrt{\pi}}\frac{v_{te}}{\omega L_{Te}} s_e \left(\frac{1}{2}Z(s_e) - s_e - s_e^2 Z(s_e)\right) \\ &+\frac{1}{4\sqrt{\pi}}\frac{v_{te}}{\omega L_{Te}}\left(1 + s_e Z(s_e)\right). \end{aligned} \tag{15}$$

The electron KTG mode is a destabilized electron sound wave with $\omega \simeq k_\parallel v_{the}$ so we retain a full plasma dispersion function in D_e to account for the electron Landau interactions. In this region $\omega/k_\parallel v_{thi} \gg 1$ so we can simplify $Z(s_i)$ functions. Solution of dispersion equation (9) with simplified response functions (12-15) and $\lambda_D = 0$ is shown in Fig. 2 by squares. We note here, that in fact plasma response is not given exactly by expressions (1-2) but contains additional terms of the order of ρ_i/L_n [18]. These additional terms do not affect asymptotic expressions (12-15) as long as $\rho_i/L_n < 1$.

A similar expansion can be made in the region $k_y\rho_i \gg 1$. Then the ion response for the ion KTG branch becomes

$$l_i = 1 - \frac{1}{2\sqrt{\pi}}\frac{v_{ti}}{\omega L_n} + \frac{1}{4\sqrt{\pi}}\frac{v_{ti}}{\omega L_{Ti}}, \tag{16}$$

$$\begin{aligned} D_i = {} & \frac{1}{2\sqrt{\pi}}\frac{v_{ti}}{\omega L_n}\left(1 + s_i Z(s_i)\right) - \frac{1}{4\sqrt{\pi}}\frac{v_{ti}}{\omega L_{Ti}} s_i \left(\frac{1}{2}Z(s_i) - s_i - s_i^2 Z(s_i)\right) \\ & - \frac{1}{2\sqrt{\pi}}\frac{v_{ti}}{\omega L_{Ti}}\left(1 + s_i Z(s_i)\right). \end{aligned} \tag{17}$$

The electron response functions can be simplified in this region by explicitly using $k_\perp\rho_e < 1$ and $\omega/k_\parallel v_{te} < 1$. Then one obtains

$$l_e = 1 - \left(1 - \frac{\omega_{ne}}{\omega}\right), \tag{18}$$

$$D_e = \left(1 - \frac{\omega_{ne}}{\omega}\right)\left(1 - 2s_e^2 + i s_e\sqrt{\pi}\right) 1 + \frac{\omega_{Te}}{\omega} s_e \left(-2s_e + i\frac{\sqrt{\pi}}{2}\right), \tag{19}$$

$$l_e = 1 - \left(1 - \frac{\omega_{ne}}{\omega}\right)(1 - b_e) - \frac{\omega_{Te}}{\omega} b_e, \tag{20}$$

$$\begin{aligned} D_e = {} & \left(1 - \frac{\omega_{ne}}{\omega}\right)\left(1 - 2s_e^2 + i s_e\sqrt{\pi}\right)(1 - b_e) + \frac{\omega_{Te}}{\omega} s_e \left(-2s_e + i\frac{\sqrt{\pi}}{2}\right)(1 - b_e) \\ & + \frac{\omega_{Te}}{\omega}\left(1 - 2s_e^2 + i s_e\sqrt{\pi}\right)\, b_e. \end{aligned} \tag{21}$$

Mode growth rate and real frequency obtained with (18-21) and (9) with $\lambda_D = 0$ are shown in Fig. 2 by circles.

Recently, it has been emphasised that the ETG modes can be significantly modified by a finite value of the Debye length parameter [19]. Electron KTG mode is strongly stabilized in a high temperature plasma by the finite Debye length as shown in Fig. 3. Note that effect of the Debye length screening may be different for the negative shear plasma [19].

Though the modified ion response for large ion Larmor radius is criticial for the ion KTG mode, the mode itself is destabilized by the electron temperature gradient as demonstrated in Fig 4. Ion KTG mode exists for large gradients of the electron temperarure (dashed line, $\eta_e = 5$). The mode growth rate is significantly reduced for smaller values of η_e (solid line, $\eta_e = 0.5$). Thus we may conclude that ion KTG mode is in fact a hybrid mode, in which both ion and electron response are essential. Electromagnetic effects are expected to be important for this mode existing in the intermediate region between standard ITG and ETG modes. Though for larger η_e we observe a standard finite β suppression (curve labeled by diamonds in Fig. 5), for low $\eta_e = 0.5$, finite plasma pressure effectively leads to the extension of the ITG instability into the shorter wavelengths (curve labeled by squares in Fig. 5). It is interesting to note that for $\beta = 0.1$ taken for this case, the skin depth size $c/\omega_{pe} \simeq 0.1\ \rho_i$, which makes $k_y c/\omega_{pe} \leq 1$ in the region of the unstable ion KTG.

In summary, we have identified two new branches of the temperature gradient driven modes. These modes are closely related to acoustic type modes that are destabilized by the temperature gradient effects. A modified plasma response in the region of $k_y \rho_\alpha \gg 1$ is essential for new instability. There is certain similarity of the KTG modes to the universal instability of drift waves that is driven by density gradient and wave-particle interaction [21]. The KTG modes considered in this work will be affected by a finite value of the magnetic shear but noting a special role of the temperature gradients and the modified plasma responce for $k_y \rho_\alpha \gg 1$ it remains unclear whether these modes can be stabilized by the shear similarly to the universal instability [22,23]. One can expect that KTG modes can persist in the region of weak and/or negative magnetic shear where the mode structure is essentially that of a shearless slab [20]. Toroidal effects will further modify the KTG modes but are not expected to cause their stabilization [24]. Generally, investigation of a general case requires nonlocal integral analysis [13,15,19] that is left for future work.

Authors would like to acknowledge useful discussions with Y. Idomura, J. Q. Li, S.I. Itoh, and K. Itoh. We are also grateful to Dr. H. Ninomiya and Dr. A. Kitsunezaki for their continuous encouragement. This research was supported by Natural Sciences and Engineering Research Council of Canada, by the Grant-in-Aid for Scientific Research and the Grant-in-Aid for International Scientific Research of Ministry of Education, Culture, Sports, Science and Technology of Japan and by collaboration program of Research Institute for Applied Mechanics of Kyushu University.

REFERENCES

[1] A. M. Dimits, G.Bateman, M. A. Beer, et al., Phys. Plasmas 7, 969 (2000).

[2] B.W. Stallard, C.M. Greenfield, G.M. Stabler et al., Phys. Plasmas **6**, 1978 (1999).

[3] Yu. Tserkovnikov, Soviet Phys. JETP **5**, 54 (1957).

[4] L.I. Rudakov and R.Z. Sagdeev, Soviet Phys Doklady **6**, 415 (1961).

[5] B.B. Kadomtsev and A.B. Timofeev, Soviet Phys Doklady **7**, 826 (1962).

[6] A.A. Galeev, V.N. Oraevskii, and R.Z. Sagdeev, Soviet Phys JETP **17**, 615 (1963).

[7] A.B. Mikhailovskii and L.I. Rudakov, Soviet Phys JETP **17,** 621 (1963).

[8] F. Jenko, W. Dorland, M. Kotchenreuthre, and B.N. Rogers, Phys Plasmas **7**, 1904 (1999).

[9] J.Y. Kim and W. Horton, Phys Plasmas B **3**, 3194 (1991).

[10] H. Biglari, P.H. Diamond, and M.N. Rosenbluth, Plasma Physics B **1**, 109 (1989).

[11] G.S. Lee and P.H. Diamond, Phys Fluids **29**, 3291 (1986).

[12] Y.C. Lee, J.Q. Dong, P.N. Guzdar, and C.S. Liu, Phys Fluids **30**, 1331 (1987).

[13] J. Q. Dong, W. Horton, and J.Y. Kim, Phys Fluids B **4**, 1867 (1992).

[14] W. Horton, B.G. Hong, and W.M. Tang, Phys Fluids **31**, 2971 (1988).

[15] J. Q. Dong, P.N. Guzdar, and Y.C. Lee, Phys. Fluids **30,** 2694 (1987).

[16] J.Y. Kim and W. Horton, Phys Fluids B **3**, 3194 (1991).

[17] Y. Chen and S. Parker, Phys. Fluids **8**, 2095 (2001).

[18] A. Smolyakov, I.O. Pogutse, and A. Hirose, Phys Plasmas **2**, 4451 (1995).

[19] Y. Idomura, M. Wakatani, and S. Tokuda, Phys. Plasmas **7**, 2456 (2000).

[20] Y. Kishimoto, J.Y. Kim, W. Horton, et al., Nuclear Fusion **40**, 667 (2000).

[21] L.D. Pearlstein and H.L. Berk, Phys. Rev. Lett **23**, 220 (1969).

[22] D.W. Ross and S. M. Mahajan, Phys. Rev. Lett **40,** 324 (1978).

[23] K.T. Tsang, P.J. Catto, J. C. Whitson, and J. Smith, Phys. Rev. Lett **40,** 327 (1978).

[24] A. Hirose, Phys Fluids B **3**, 1125 (1991).

FIGURES

FIG.1. Normalized (to the ion cyclotron frequency) frequency (a) and growth rate (b) of the ITG and unstable drift wave modes versus the normalized ion Larmor radius ρ_i/L_n. The solid line corresponds to the ITG branch, while the dashed line is that of the destabilized drift wave branch. Parameters are the same as those in Ref. $\beta = 2 \times 10^{-4}$, $\rho_i/L_{Ti} = \sqrt{2} \times 10^{-1}$, $\rho_i/L_{Te} = \sqrt{2} \times 10^{-2}$, $k_x\rho_i = \sqrt{2} \times 10^{-1}$, $k_y\rho_i = 3\sqrt{2} \times 10^{-1}$, $k_\parallel\rho_i = 2\sqrt{2} \times 10^{-3}$.

FIG.2. Normalized wave frequency (a) and growth rate (b) for the ITG , ETG modes and kinetic modes. Plasma parameters: $\beta = 2 \times 10^{-4}$, $\rho_i/L_n = \sqrt{2} \times 10^{-2}$, $\rho_i/L_{Ti} = \sqrt{2} \times 10^{-1}$, $\rho_i/L_{Te} = \sqrt{2} \times 10^{-1}$, $k_x\rho_i = \sqrt{2} \times 10^{-1}$, $k_\parallel\rho_i = 2\sqrt{2} \times 10^{-3}$, $\tau = 1$. Solid line – standard ITG and ion KTG; dashed line – standard ETG and electron KTG.

FIG.3. Stabilization of the Kinetic Electron Temperature Gradient mode by the Debay screening effect: solid line – $\lambda_D/\rho_e = 0$, dotted-dashed line – $\lambda_D/\rho_e = 0.70$, dotted line – $\lambda_D/\rho_e = 2.21$. Plasma parameters: $\beta = 2 \times 10^{-4}$, $\rho_i/L_n = \sqrt{2} \times 10^{-2}$, $\rho_i/L_{Ti} = \sqrt{2} \times 10^{-1}$, $\rho_i/L_{Te} = \sqrt{2} \times 10^{-1}$, $k_x\rho_i = \sqrt{2} \times 10^{-1}$, $k_\parallel\rho_i = 2\sqrt{2} \times 10^{-3}$, $\tau = 1$.

FIG.4. Effect of the electron temperature gradient on KTG modes. Solid line – $\rho_i/L_{Te} = \sqrt{2} \times 10^{-2}$ ($\eta_e = 0.5$), dashed line $\rho_i/L_{Te} = \sqrt{2} \times 10^{-1}$ ($\eta_e = 5$). Plasma parameters: $\beta = 2 \times 10^{-4}$, $\rho_i/L_n = \sqrt{2} \times 10^{-2}$, $\rho_i/L_{Ti} = \sqrt{2} \times 10^{-1}$, $k_x\rho_i = \sqrt{2} \times 10^{-1}$, $k_\parallel\rho_i = 2\sqrt{2} \times 10^{-3}$, $\tau = 1$.

FIG.5. Effect of a finite plasma pressure on the KTG modes. Triangles – $\rho_i/L_{Te} = \sqrt{2} \times 10^{-1}$, $\beta = 2 \times 10^{-4}$; diamonds squares – $\rho_i/L_{Te} = \sqrt{2} \times 10^{-1}$, $\beta = 0.1$; circles $\rho_i/L_{Te} = \sqrt{2} \times 10^{-2}$, $\beta = 2 \times 10^{-4}$, squares – $\rho_i/L_{Te} = \sqrt{2} \times 10^{-2}$, $\beta = 0.1$. Other plasma parameters: – $\rho_i/L_n = \sqrt{2} \times 10^{-2}$, $\rho_i/L_{Ti} = \sqrt{2} \times 10^{-1}$, $k_x\rho_i = \sqrt{2} \times 10^{-1}$, $k_\parallel\rho_i = 2\sqrt{2} \times 10^{-3}$, $\tau = 1$.



Fig.1a   A. I. Smolyakov 10/23/01



Fig.1b A. I. Smolyakov 10/23/01



Fig.2a   A. I. Smolyakov 10/23/01



Fig.2b A. I. Smolyakov 10/23/01



Fig.3 A. I. Smolyakov 10/02/01



Fig.4 A. I. Smolyakov 10/02/01



Fig.5 A. I. Smolyakov 10/02/01